

Ph: 478 742 8766 Website: Hudspethanimalhospital.com

## **Patient Drop Off Form**

The information provided will allow our team to provide the care your pet needs. It is our pleasure to provide superior care.

Today's Date:/	_/	_		
Pet's Name: Breed:				
Color:		Sex:		Age
We will need to contact you operson responsible? Me of		•		edical and financial decisions. Who is the
Phone numbers of responsible	le persons.			
First phone				
Second phone				
Reason(s) for Drop o	off			
Vaccination		Bath		Illness
Bi-annual Comprehensive		Haircut		Vaccine Allergy Treatment
Examination		Training		Microchipping
		Bandage Change		
Dental Prophylaxis	Ш	bandage change		

Is your pet taking any medication(s)? ☐ Yes NO ☐						
Name of Medication	Dose	Dat	e and time Last Given	Reason of medication		
Has your pet ever had an adve	rse reaction to vac	cines or	any medications? 🔲	∕es □No		
	Name o	of Medica	tion			
Please identify area(s) of co	ncern (Circle con	cerns):				
Eating Drinking	Bad Breath		Excessive Sleeping			
Weight Loss	Weight Gain		Itching/Scratching			
Difficulty Rising	Scooting		Shaking Head			
Vomiting Diarrhea	Masses/Lesions		Urination Is	Urination Issues		
Behavioral Problem	eyes		Skin			
Other						
Please call me before treating i	f my fee will be ove	er \$	(if I	eft blank, we will call if		
fee is over \$100) or If fee will b	e \$					

Preventative Services Offered – Indicate what service you would like today (additional				
charges apply).				
<b>Pre-med Prior to Vaccinations:</b> Hudspeth Animal Hospital takes step possible to minimize the				
chance of an allergic reaction to vaccines. Give an antihistamine 20 – 30 minutes before				
vaccination reduces, however, does not eliminate the risk of vaccine allergic reactions. This is				
recommended in small dogs and breeds prone to vaccine allergic reactions. A diphenhydramine				
injection prior to vaccination, or after vaccination to lessen its severity if they do react.				
Additional medications or treatments may be needed for severe vaccine allergic reactions.				
<b>Heartworm Prevention</b> : Special worms cause heartworm disease. The worms are called				
Dirofilaria immitis and are transmitted by mosquitoes. The worms live the heart and big vessel of				
the heart. They often migrate into the lungs. These worms cause irreversible damage to the				
heart leading to heart disease. Infected dogs will die without appropriate treatment. Cats are				
also capable of getting heartworm disease. Signs of infection in cats includes sudden death,				
vomiting, coughing, respiratory distress. Heartworm disease is 100 % preventable by giving				
appropriate heartworm preventative to cats and dogs. Appropriate heartworm testing is				
required before prevention may be administered in dogs. Heartworm testing is also				
recommended in cats.				
Anal Gland Expression: Anal sacs are found along the edge of the anus. They may get blocked,				
infected and full. They normally empty when the pet has a bowel movement, however, if the				
secretion is thick, it will not be able to flow out. Signs of issue with the anal sacs may include				
licking, chewing, scooting, discharge around the anus, and pain. Manual expression will become				
necessary if blocked.				
Flea and Tick Prevention: Tick, fleas, roundworms, and hookworms are common parasites of				
companion animals that can cause disease in humans. Keeping your pets on preventatives				
protects both pet and owner. We can help you choose the appropriate heartworm preventative				
for your pet. A flea control product will be administered to pets with fleas. This will be done at				
the owner's expense. This is to protect animals in the hospital and prevent infestation of the				
hospital.				

As owner / agent / caregiver I have read, understand and agree to the aforementioned.

I agree and understand that as caregiver / owner / agent I am responsible for all charges incurred.

Print Name:	Date:
Signaturo	